

Upper Missouri River Water Quality Report 2021: from Lewis & Clark, Broadwater and Cascade Counties, MT

Prepared for:

Sherry Meador, Alan Shaw, Bailey Sory and Pat Hunter
Project Managers/Co-Chairs/Board
Upper Missouri Watershed Alliance UMOWA
Wolf Creek, Montana



Water quality collection station at the Missouri River looking upstream of Little Prickly Pear Creek

Prepared by:

Dave Stagliano,
Montana Biological Survey
February 2022



Table of Contents

Executive Summary.....	3
1.0 Introduction	5
2.0 Methods.....	6
2.1 Sample Site Selection.....	6
2.2 Sample Timing.....	7
2.3 Water Sampling Methods.....	8
2.4 Quality Assurance – Field Blanks and Duplicates	9
2.5 Water Quality Analysis.....	9
3.0 Results for 2021	9
3.1 Total Dissolved Solids.....	10
3.2 Total Suspended Solids	11
3.3 Inorganic Nitrogen	12
3.4 Total Nitrogen	13
3.5 Total Phosphorus	12
4.0 Discussion.....	15
4.1 Spatial Trends.....	15
4.2 Seasonal Trends	15
4.3 Yearly Trends.....	16
5.0 Conclusion.....	16
6.0 References	17
Appendix A: Yearly water quality data for 2016-2021.....	18

Acknowledgements

We would like to thank Alan Shaw, Sherry Meador, Pat Hunter, Bailey Sory and the rest of the UMOWA Board for initiating and continued support for this important project. We thank Braden Lewis for collecting the WQ samples for the first 3 years and preparing previous reports. Montana DEQ has provided funding for the water sample analysis through Energy Laboratories, Inc. in Helena, MT. Katie Makarowski and Abbie Ebert, MDEQ Watershed Planners were especially helpful in developing our WQ SAP as well as navigating us through the 319-watershed grant funding process. Deanna Tarum has been especially helpful in getting our data into DEQ’s MT-eWQX EQUIS database

Executive Summary

In 2016, the Upper Missouri Watershed Alliance (UMOWA) began collecting water quality (WQ) data at seven sites on the Upper Missouri River between Wolf Creek and Cascade. These sites were chosen to coincide with long-term macroinvertebrate monitoring sites that were established in 2015 by Montana Biological Survey. 2021 marks the 6th consecutive year that WQ samples have been collected. WQ samples were usually collected seasonally in April/May, July and late-September/October to coincide with pre-, post- and during irrigation periods.

The goals of the project are: 1) to continue to collect seasonal baseline WQ data as it may affect benthic macroinvertebrate, algae and plant communities and 2) to collect baseline information for total suspended solids (TSS), total dissolved solids (TDS) and nutrients (Total Nitrogen, TN; inorganic nitrogen, NN; total phosphorus, TP) to allow for spatially and temporal robust comparisons of water quality in the Upper Missouri River. WQ samples have not been analyzed for recoverable metals since 2016, because then, these metals were recovered in low (often non-detectable) levels (except an Arsenic (AS) spike during spring below Little Prickly Pear Creek) suggesting that they pose, little to no acute or chronic threat to the Missouri River.

The water quality data collected in 2021 has reinforced some of the seasonal and spatial trends observed since 2016 but has also indicated a reverse of the increasing trend of TN and NN concentration levels in the Missouri River below Holter. Additional tributary sites sampled for WQ upstream of

Canyon Ferry Reservoir revealed some very high levels of TP and TN entering the Missouri River upstream of Townsend.

During the fall sampling period of 2021, TN concentration levels at two sites and TP levels at 5 of the 7 monitoring sites below Holter Dam exceeded the old, recommended standards set by MDEQ. Overall monitoring years, nutrient concentrations (TN and TP) were the highest in 2018 compared to 2016, 2017, 2019-2021. This increase was most apparent during June and October. Inorganic nitrogen (NN) concentrations have remained at very low concentrations below Holter since 2019, and the MDEQ numeric nutrient standards for TN, and (with MO Cascade) for TP.

Nutrient levels in 2021, particularly total phosphorus (TP), were lower downstream of Holter Dam than in previous years. Lower mean discharge and no flushing flows in 2021 did not provide the lower Missouri River below Canyon Ferry with a large nutrient pulse that occurred in 2017-2020. The Missouri River upstream of Canyon Ferry exceeded the old MDEQ nutrient thresholds for both Total Nitrogen (TN) and TP in July, but these nutrients are being utilized within the reservoir. The tributaries contributing these excess nutrients are Big Springs Ditch (TN), Deep Creek (TP) and the biggest contributor was Greyson Creek for both TN and TP with over 5 times the TP 'allowed' by MDEQ. Nutrient levels in the mainstem Missouri River have been trending upward over the 4 study years, most notably in the Spring and Fall samples, while below tributaries, concentrations have been mediated by

the inflows; this rising trend has stabilized in 2020. October 2017 was really at the forefront of the increasing trends in nutrient concentrations during the previous sampling periods.

It is evident that high nutrient loads entering Canyon Ferry are the biggest threat that the Upper Missouri River is facing at present. Zero sites sampled in 2016 during any season exceeded the old numeric nutrient standards or screening values set by MDEQ, but now most monitoring sites are surpassing these thresholds routinely in the fall (UMOWA 2020). It is critical that UMOWA continue to monitor the Missouri River water quality into 2022 to see if nutrient concentrations continue to stabilize below impairment levels or continue an upward trajectory.

1.0 Introduction

The 2.8-mile reach of the Missouri River from Holter Dam to Little Prickly Pear Creek is listed as impaired for aquatic life by Montana Department of Environmental Quality (MDEQ) due to sedimentation/siltation, increased total nitrogen and phosphorus concentrations (likely caused by grazing in riparian zones, municipal point source discharges and caused by on-site treatment systems (septic systems) and flow regime alterations from upstream impoundments (MDEQ 2016). The 20.9-mile reach of the Missouri River from Little Prickly Pear Creek to Sheep Creek is listed as impaired for aquatic life due to increased arsenic concentrations, increased total nitrogen concentrations due to grazing in riparian zones, impacts from hydro-structure flow regulation/modification, and sedimentation/siltation caused by irrigated crop production and natural sources (USEPA 2020). Similar impairments are listed for the next Missouri River section from Sheep Creek to the Sun River (USEPA 2020). Nutrient levels in the mainstem Missouri River have been trending upward over the past 4 study years, most notably in the Spring and Fall samples, while below tributaries, concentrations are somewhat mediated by the inflows (UMOWA 2021).

In 2016, the Aquatic Resources Committee of the Upper Missouri Watershed Alliance (UMOWA) decided to fund the collection of water quality data on a periodic and consistent basis to monitor the health of the Upper Missouri River between Holter Dam and Cascade. Samples were collected concurrently with macroinvertebrate samples being collected by Montana Biological Survey (Stagliano 2019) to determine if any correlations can be drawn between water chemistry and aquatic insect communities. The overall goals of the project were to 1) establish baseline water quality data as it may affect benthic macroinvertebrate and aquatic plant communities, and 2) collect baseline information about total suspended/dissolved solids and nutrients to allow for spatial and temporal comparisons of water quality. To get an idea of river assessment conditions, WQ data will be compared against nutrient standards for wadable streams in the same ecoregion. In 2017, analysis of metal concentrations was eliminated from the project goals. The 2018-2021 WQ sampling focused on nutrient concentrations as well as total suspended/dissolved solids concentrations.

From our 6 years of collecting and analyzing spatial and temporal WQ data, it has become clear that most of the nutrient issues on the Missouri River below Holter Dam are not from local sources but are originating from one or more of the upstream reservoirs and must be addressed at their sources. Therefore, in 2021 we sampled WQ nutrient data at 8 new Missouri River or Canyon Ferry tributaries that are on the 303(d) Impaired Waters List with nutrients: TN, TP and/or NN, listed as at least one cause of Aquatic Life impairment (MDEQ Appendix A 2020).

2.0 Methods

2.1 Sample Site Selection

Water quality samples were collected seasonally from seven sites between Wolf Creek and the town of Cascade (Table 1, Figure 1). These sites coincide with monitoring sites that have been sampled for macroinvertebrates by Montana Biological Survey since 2015. The sites were chosen to be spatially representative of the upper Missouri River and integrate the influence of the three major tributaries (Little Prickly Pear Creek, Dearborn River and Sheep Creek) on water quality.

Station ID	Site Name	Latitude	Longitude
MO_LPPC_US	Missouri River U/S Little Prickly Pear	47.02281	-112.01527
MO_LPPC_DS	Missouri River D/S Little Prickly Pear	47.02345	-112.01523
MO_CRAIG	Missouri River U/S Craig	47.05415	-111.96701
MO_DEAR_US	Missouri River U/S Dearborn River	47.12819	-111.91174
MO_DEAR_DS	Missouri River D/S Dearborn River	47.12791	-111.9109
MO_HARDY	Missouri River U/S Sheep Creek (Hardy Bridge: cover photo)	47.16781	-111.83366
MO_CASCADE	Missouri River @ Cascade FAS	47.28062	-111.69113

Figure 1. WQ Sampling sites are denoted by blue triangles. Black dots are towns.





Table 2. Tributary stream locations proposed for WQ Sampling in 2021.

Station ID	Agency	Site Name	Latitude	Longitude	Parameters to Collect**
MT41I002_030	UMOWA	Beaver Creek, headwaters to mouth, sample D/S of HWY 12	46.5134	-111.5995	WQS (dry, NS*)
MOtrib_BigSpring	UMOWA	Big Springs Ditch U/S of HWY12	46.18146	-111.4447	WQS (added)
MT41I002_042	UMOWA	Confederate Gulch, Hunter Gulch to mouth (Canyon Ferry Reservoir)	46.5003	-111.5106	WQS (dry, NS*)
MT41I002_050	UMOWA	Crow Creek (USFS boundary to mouth of Missouri River)	46.1874	-111.4843	WQS, NS*
MO trib_DeepCR	UMOWA	Deep Creek D/S of HWY 12	46.28603	-111.4996	WQS (added)
MT41I002_080	UMOWA	Dry Creek, National Forest boundary to Missouri River	46.2436	-111.4509	WQS
MO trib_Greyson	UMOWA	Greyson Creek D/S of HWY 12	46.26329	-111.4842	WQS (added)
MT41I002_110	UMOWA	Magpie Creek, headwaters to Canyon Ferry Reservoir	46.6449	-111.6809	WQS
MO_Townsend	UMOWA	Missouri River bridge in Townsend	46.33527	-111.5325	WQS (added)
MT41I002_120	UMOWA	Sixteenmile Creek, Lost Creek to Missouri River	46.1066	-111.3966	WQS
MT41I002_150	UMOWA	Cave Gulch, headwaters to mouth (Canyon Ferry Reservoir)	46.6611	-111.6971	WQS (dry, NS*)
MT41I006_080	UMOWA	Spring Creek, Corbin Creek to mouth (Prickly Pear Creek)	46.3885	-112.0306	WQS

**WQS=Water quality samples (TSS, TDS, NO2+3, TP, TN), *NS-not sampled


Map 2. WQ Sampling sites upstream of Holter Reservoir are denoted by blue drops.



2.2 Sample Timing

Samples were collected seasonally to examine temporal changes in water quality through the year, especially related to the summer irrigation/growing season. In 2021, seasonal samples were collected on May 15th, July 20th & 22nd and September 30st, similar dates as in 2020, while in 2019, only a fall sample was collected (October 4th) (**Figure 2**). WQ samples collected in 2021 captured data during some of the lowest annual discharge reported in the last decade; there wasn't even a spring pulse (**Figure 2**). Three proposed tributary sites were dry when visited in July.

Figure 2. Discharge below Holter Dam from Jan 2019-2021. Red arrows represent sampling dates



2.3 Sampling Methods

Water quality samples were collected in sterilized bottles provided by Energy Laboratories Inc. At the time of sample collection, water bottles were held upstream of the collector and each bottle and lid were triple rinsed with ambient river water prior to collecting the final sample. Total nitrogen (TN) samples were collected in 250 mL HDPE bottles and kept on ice until analyzed. Total phosphorus (TP) and inorganic nitrogen (nitrate and nitrite, NN) samples were collected in 250 mL HDPE bottles, preserved with sulfuric acid and nitric acid, respectively, and kept on ice until analyzed. Total suspended solids (TSS) and total dissolved solids (TDS) samples were

collected in 1000 mL HDPE bottles and kept on ice until analyzed. All water samples were submitted to Energy Laboratories Inc. in Helena, MT within 24 hours of collection.

2.4 Quality Assurance – Field Blanks and Duplicates

During each seasonal sampling, one field duplicate and one field blank set of samples were collected to ensure the quality of the sampling procedure and the accuracy of the lab analysis. Field duplicates consist of an extra set of sample bottles filled with the same water from the same sampling site. Field duplicates were collected, handled, and stored in the same way as the routine samples for laboratory shipment. In theory, a duplicate sample should yield results identical to the other sample taken at the same site at the same time. Collecting and analyzing duplicate samples ensures that the sampling and analysis process is consistently accurate.

Field blanks consist of an extra set of sample bottles filled with deionized (DI) water and transported with the other samples in the field. Field blanks were prepared in the field at the same time as the routine samples and handled and analyzed in the same way as the routine samples. In theory, field blank samples should be clean of all contaminants. Collecting and analyzing field blank samples ensures that the sampling and analysis processes are free of contaminants.

2.5 Water Quality Analysis

Energy Laboratories Inc. in Helena, MT analyzed all water quality samples. Energy Laboratories Inc. routinely processes 1,000's of WQ samples for MDEQ, USEPA and other agencies, and has been proven to provide laboratory data of very high accuracy and consistency.

3.0 Results for 2021

Three tributary sites proposed for sampling this year (Cave Gulch, Confederate Gulch and Beaver Creek) were dry when visited in late-July; therefore, we sampled Big Springs Ditch, Deep and Greyson Creeks, and the Missouri River in Townsend as alternatives (**Table 2, Map 2**). Whenever possible, results were interpreted relative to previously established MDEQ numeric WQ standards within the ecoregion. MDEQ established these numeric water quality criteria to control limits on pollutant levels in Montana's wadable rivers and streams. Numeric nutrient criteria had not yet been developed for the Missouri River (a non-wadable river), and wadable stream nutrient thresholds have been discontinued in 2021 in favor of narrative criteria. Circular 12A (MDEQ 2014) provides the discontinued threshold numeric nutrient standards of 0.3 mg/L for total nitrogen (TN) concentrations and 0.03 mg/L for total phosphorus (TP) concentrations. The screening value of 0.1 mg/L for inorganic nitrogen concentrations was provided via personal communication with MDEQ staff. Screening values and numeric nutrient standards are represented as horizontal lines on the graphs. A table of all 2016-2021 WQ results is available in Appendix A.

3.1 Total Dissolved Solids

In 2021, Total Dissolved Solid (TDS) concentrations ranged from 143 and 508 mg/L and averaged 240 mg/L across all sites and seasons; no distinct spatial trends were noted below Holter Dam except lower May concentrations at the Missouri River below Little Prickly Pear Creek (MO_LLPC_DS) (**Figure 3**). Seasonally, July 2021 samples had the highest TDS levels downstream of the Dearborn and May samples had the lowest TDS, except at the Missouri sites closer to the Dam. located below tributaries where fall reported the highest levels (**Figure 3**). In 2019, TDS concentrations ranged between 203 and 217 mg/L and averaged 209 mg/L across all sites; again, the highest levels were reported at sites below the major tributaries in the Fall (**Figure 3**).







Figure 3. Total Dissolved Solid levels across the UMOWA sites in 2021 and 2017-2021. Sites arranged upstream to downstream

3.2 Total Suspended Solids

In 2021, Total suspended solids (TSS) concentrations ranged between 1.5 and 104 mg/L and averaged 10.2 mg/L across all sites and seasons; downstream of Holter Reservoir, average TSS values were 4.6 mg/l across all sites and seasons (**Figure 4**). Not surprisingly, the highest TSS values reported downstream of Holter in 2021 and in the entire study period, were reported from the samples taken downstream of Little Prickly Pear Creek (26mg/l), the Dearborn River and Sheep Creek (Cascade) during those streams’ high-flow May period (**Figure 4**). Most of the tributaries upstream of Canyon Ferry had substantially higher TSS values than below Holter. The Missouri River at Hardy Bridge had non-detectable TSS levels during all seasons in 2020 and from July 2016 to Oct 2017, but this year had more detectable levels (**Figure 4**).

Figure 4. Total Suspended Solids (TSS) across the UMOWA sites in 2021 (top) and 2016-2020 (bottom). Sites arranged upstream to downstream






3.3 Inorganic Nitrogen

In 2021, Total Inorganic Nitrogen (NN) concentrations ranged between 0.0 and 0.25 mg/L and averaged 0.04 mg/L across all sites and seasons: with zero detectable levels downstream of Holter Dam in July and one detectable sample in September at MO_LPPC_US. No sites below Holter approached the MDEQ screening value in May, while upstream 3 values exceeded this level in July (Figure 5). The highest NN values between 2016 and 2020 were reported from samples taken during the Oct. 2018 sampling period and lowest during the May 2020 period where all sites except below Little Prickly Pear and all seasons at Cascade in 2020 reported non-detectable levels (Figure 5).



Figure 5. Total Inorganic Nitrogen (NN) levels across the UMOWA sites in 2021 (top) and 2016-2021 (bottom). Sites arranged upstream to downstream



3.4 Total Nitrogen

In 2021, Total Nitrogen (TN) concentrations ranged between 0.07 and 0.57 mg/L and averaged 0.27 mg/L (± 0.03 SE) across all sites and seasons; only 2 sites below Holter were above the MDEQ old nutrient standard for wadable streams in October and one in May, while upstream 5 of the 8 sites exceeded the 0.3 mg/l threshold in July (Figure 6). Elevated TN concentrations from the Dry Creek and Greyson Creek tributaries increased the TN concentrations at the Missouri River near Townsend to levels above the old MDEQ standard (Figure 6). October 2017 was the 1st sample period where Missouri River samples began exhibiting elevated TN levels (Figure 6).

Figure 6. Total Nitrogen (TN) levels across the UMOWA sites in 2021 (top) and 2016-2021 (bottom). Sites arranged upstream to downstream



3.5 Total Phosphorus

In 2021, Total phosphorus (TP) concentrations ranged between 0.012 and 0.157 mg/L and averaged 0.032 mg/L across all sites and seasons; 5 of the 8 sites sampled in July (upstream of Holter) and 5 of the 7 sites sampled in late-September (d/s Holter Reservoir), reported values above the MDEQ old nutrient standard threshold (**Figure 8**). Greyson Creek reported very elevated TP levels (5 times the old nutrient standard). Elevated TP concentrations were also reported below Holter during October 2018, 2019 and 2020 with 5 or 6 of the 7 total sample sites reporting levels above the MDEQ old standard (**Figure 9**).

Figure 8. Total Phosphorus levels across the UMOWA sites in 2021.





Figure 9. Total Phosphorus levels across the UMOWA sites during 2016-2021.



4.0 Discussion

4.1 Spatial Trends

When comparing Missouri River upstream sites closer to Holter Dam to downstream sites, no strong trends were observed for any of the measured WQ parameters in 2021, except below the tributaries where concentrations tended to be diluted. When nutrients were exceeding the MDEQ old numeric nutrient thresholds, they are generally elevated from upstream of Little Prickly Pear Creek all the way downstream to Cascade, usually with only slight decreases in the concentration levels (**Figures 5, 6 & 8**). There were no documented “big spikes or dips” between sites unless a tributary comes in and dilutes levels down for a kilometer or 2 km until full mixing occurs, then nutrient levels go right back up. There were a couple of notable exceptions; the Missouri River samples taken below Little Prickly Pear Creek and the Dearborn River reported reduced nutrient levels below MDEQ nutrient standards in all seasons, except June 2018 and Oct 2017. During October 2016, 2018 and 2019, the MO_DEAR_DS site yielded exceptionally low and/or non-detectable TP concentrations (**Figure 10**). Although, concentrations of all three nutrient parameters (NN, TN and TP) are increasing in the Missouri River below the Dearborn site (MO_DEAR_DS) since 2019 indicating that lower Dearborn River flows are offering less of a dilution effect. The high TN and TP concentrations that we documented in some of the tributaries upstream of Canyon Ferry Reservoir are elevating the readings in the Missouri River but are likely being utilized with the reservoir and are not continuing to spiral downstream below Holter.

4.2 Seasonal Trends

Total dissolved solids (TDS) concentrations were fairly uniform from May through September for six of the seven sites in 2021; only the Missouri River downstream of Little Prickly Pear Creek had lower values in the spring than reported in the summer or fall. This may be due to early run-off from LPPC and some dilution effects, but was not seen below the Dearborn River. The Missouri River below LPPC reported a high spike in TSS in May of 2020. It is unclear as to what extent natural and anthropogenic sources (fertilizers, pesticides, etc.) are contributing to elevated TDS concentrations during the June run-off period (Total Nitrogen is also very elevated during this period). But high readings of TDS were also reported at all sites in Fall of 2019, especially below the tributaries.

Overall, in 2021, nutrient levels were greater in September than in May or July; TP was significantly higher in October than May (T-test, $p=0.001$) or July (T-test, $p=0.025$). Total nitrogen, and total phosphorus concentrations increased between July and September for six of the seven sites (the one exception was the TP levels at MO_CRAIG as described in 4.1).

4.3 Yearly Trends

Total dissolved solids (TDS) concentrations have been higher in 2021 and 2020 than reported in 2016-2019, although, this parameter was inconsistently sampled. This could be related to lower discharge and the concentrating effect of lower river volumes (**Figure 2, Appendix A**). TSS concentrations reached their highest levels in April 2016, June 2018 and May 2020, but they were

similar between Fall 2018 and 2019. TSS concentrations were zero (i.e., at non detectable levels) at all sites between July 2016 and October 2017, except at the Cascade site, which exhibited moderate concentrations during July of all years. TSS concentrations has been correlated with or a contributor to higher nutrient levels.

5.0 Conclusions

This study has established a comprehensive baseline of the water quality in the upper Missouri River over six different water discharge years (2016-2021) and has elucidated several spatial and temporal trends. The 2021 water quality downstream of Holter Reservoir has been trending towards lower nutrient concentrations (TN, TP, and NN) across all seasons compared to October 2017-2019 when the overwhelming trends observed were substantial increases in inorganic nitrogen (NN), total nitrogen (TN), and total phosphorus (TP) concentrations to levels exceeding MDEQ old nutrient standards (UMOWA 2020). Nutrient exceedances in 2018 and 2019 were reported from the furthest upstream site all the way down to Cascade, except at sites below the tributaries, Little Prickly Pear Creek and the Dearborn River were somewhat insulated from these increasing nutrient trends, but some seasonal increases are being documented.

During September 2021 sampling, two and five of the seven sites exceeded the MDEQ old nutrient standard values for TN and TP concentrations, respectively. Zero sites exceeded the MDEQ recommended screening value for NN. It appears that TN levels have declined in 2020 and 2021 after nutrient concentrations exceeding the MDEQ nutrient standards and screening values, across most sites, have occurred during the Fall sampling of 2017, 2018 and 2019. Warmer September temperatures and lower discharge in 2020 and 2021 with numerous sunny days may be one explanation for lower TN levels, as a longer growing season means that aquatic plants would be utilizing more nutrients than they would if water temperatures had dropped earlier in autumn.

Although we have observed fewer nutrient levels (TN, NN) with concentrations exceeding the recommended levels set by MDEQ in 2021, high TP levels at 5 of 7 sites in the fall of 2021 is still a cause for concern. Higher nutrient levels are causally related to the excessive aquatic plant and algae growth are among the biggest threats that the Upper Missouri River is currently facing. In terms of nutrients and aquatic plant growth, weed growth in 2021 may not have exploded as one might have expected with the warm temperatures. Due to UMOWA's water quality collections, we have a pretty good explanation for why this is the case; nutrient levels, particularly total phosphorus (TP), were lower downstream of Holter Dam than in previous years. Lower mean discharge and no flushing flows in 2021 did not provide the lower Missouri River below Canyon Ferry with a large nutrient pulse as occurred in 2017-2020. The Missouri River upstream of Canyon Ferry still exceeded the old MDEQ nutrient thresholds for both Total Nitrogen (TN) and TP in July, but these nutrients are being utilized within the reservoir. The tributaries contributing these excess nutrients are Big Springs Ditch (TN), Deep Creek (TP) and the biggest contributor was Greyson Creek for both TN and TP with over 5 times the TP 'allowed' by MDEQ. It is critical that UMOWA continue to monitor water quality parameters in 2021 to see if these nutrient

concentrations continue to reach impaired levels. Additional upstream watershed BMP measures (riparian vegetation buffers, cattle fencing, etc.) should be evaluated to alleviate non-point source nutrient additions to the ecosystem.

6.0 References

MDEQ 2020. Appendix A: Impaired Waters. (2020). Available at:
[http://deq.mt.gov/Portals/Impaired Waters2020](http://deq.mt.gov/Portals/Impaired_Waters2020)

Montana DEQ 2014. Planning Prevention and Assistance Division, Water Quality Planning Bureau, Water Quality Standards Section. 2014. DEQ Circular-12A Montana Base Numeric Nutrient Standards. Helena, MT: Montana Dept. of Environmental Quality.

Stagliano, D. 2015. UMOWA Baseline Macroinvertebrate Monitoring Report 2015 for the Upper Missouri River, MT. Report to the Upper Missouri River Alliance.

Stagliano, D. 2018. UMOWA Baseline Summary Macroinvertebrate Monitoring Report for 2015-2017 for the Upper Missouri River, MT. Report to the Upper Missouri River Alliance.

UMOWA. 2021. Baseline Water Quality Report 2020 for the Upper Missouri River, MT. Report to the Upper Missouri River Alliance prepared by D. Stagliano.

APPENDIX A. WATER QUALITY RAW DATA FROM 2016-2021

Sample Site	Year	Sample Time	TDS (mg/L)	TSS (mg/L)	Inorganic Nitrogen (mg/L)	Total Nitrogen (mg/L)	Total Phosphorus (mg/L)
MO_LPPC_US	2016	April	NA	0	0	0.24	0.014
MO_LPPC_DS	2016	April	NA	14	0.05	0.23	0.02
MO_Craig	2016	April	NA	4	0	0.22	0.016
MO_Dearborn US	2016	April	NA	4	0	0.22	0.018
MO_Dearborn DS	2016	April	NA	7	0	0.09	0.01
MO_Hardy Creek	2016	April	NA	5	0	0.23	0.021
MO_Cascade	2016	April	NA	8	0	0.26	0.019
MO_LPPC_US	2016	July	NA	0	0	0.13	0.011
MO_LPPC_DS	2016	July	NA	0	0	0.04	0.008
MO_Craig	2016	July	NA	0	0	0.15	0.012
MO_Dearborn US	2016	July	NA	0	0	0.15	0.012
MO_Dearborn DS	2016	July	NA	0	0	0.14	0.013
MO_Hardy Creek	2016	July	NA	0	0	0.14	0.013
MO_Cascade	2016	July	NA	5	0	0.14	0.015
MO_LPPC_US	2016	October	NA	0	0.07	0.28	0.023
MO_LPPC_DS	2016	October	NA	0	0.02	0.18	0.017
MO_Craig	2016	October	NA	0	0.06	0.26	0.02
MO_Dearborn US	2016	October	NA	0	0.06	0.26	0.019
MO_Dearborn DS	2016	October	NA	0	0	0.06	0
MO_Hardy Creek	2016	October	NA	0	0.05	0.25	0.017
MO_Cascade	2016	October	NA	0	0	0.19	0.015
MO_LPPC US	2017	July	216	0	0	0.18	0.013
MO_LPPC DS	2017	July	197	0	0.04	0.12	0.016
MO_Craig	2017	July	207	0	0	0.22	0.012
MO_Dearborn US	2017	July	202	0	0.05	0.22	0.012
MO_Dearborn DS	2017	July	187	0	0	0.09	0.004
MO_Hardy Creek	2017	July	208	0	0	0.2	0.012
MO_Cascade	2017	July	206	4	0	0.21	0.014
MO_LPPC US	2017	October	197	0	0.14	0.38	0.068
MO_LPPC DS	2017	October	203	0	0	0.11	0.016
MO_Craig	2017	October	195	0	0.14	0.36	0.068
MO_Dearborn US	2017	October	194	0	0.15	0.37	0.067
MO_Dearborn DS	2017	October	191	0	0.15	0.36	0.069
MO_Hardy Creek	2017	October	190	0	0.15	0.34	0.063
MO_Cascade	2017	October	193	0	0.08	0.29	0.055

Sample Site	Year	Sample Time	TDS (mg/L)	TSS (mg/L)	Inorganic Nitrogen (mg/L)	Total Nitrogen (mg/L)	Total Phosphorus (mg/L)
MO_LPPC US	2018	June	203	6	0.14	0.45	0.031
MO_LPPC DS	2018	June	202	2.4	0.14	0.43	0.03
MO_Craig	2018	June	206	10	0.14	0.44	0.037
MO_Dearborn US	2018	June	204	6	0.15	0.44	0.033
MO_Dearborn DS	2018	June	166	6	0	0.08	0.006
MO_Hardy Creek	2018	June	204	7	0.14	0.41	0.028
MO_Cascade	2018	June	206	9	0.14	0.42	0.035
MO_LPPC US	2018	July	177	1.2	0.08	0.38	0.033
MO_LPPC DS	2018	July	198	2.8	0.06	0.24	0.018
MO_Craig	2018	July	173	1.6	0.03	0.32	0.031
MO_Dearborn US	2018	July	163	1.6	0.04	0.32	0.028
MO_Dearborn DS	2018	July	192	0.8	0	0.11	0.006
MO_Hardy Creek	2018	July	167	1.2	0.03	0.33	0.027
MO_Cascade	2018	July	168	4	0	0.3	0.025
MO_LPPC US	2018	October	179	2.4	0.17	0.49	0.08
MO_LPPC DS	2018	October	208	2.4	0.07	0.24	0.017
MO_Craig	2018	October	181	2.8	0.17	0.48	0.081
MO_Dearborn US	2018	October	177	1.2	0.17	0.46	0.079
MO_Dearborn DS	2018	October	205	0.8	0	0.07	0
MO_Hardy Creek	2018	October	179	2	0.19	0.47	0.078
MO_Cascade	2018	October	180	1.2	0.14	0.47	0.076
MO_LPPC US	2019	October	206	2.8	0.11	0.34	0.06
MO_LPPC DS	2019	October	220	1.6	0.1	0.21	0.033
MO_Craig	2019	October	209	2	0.12	0.29	0.056
MO_Dearborn US	2019	October	207	0.8	0.09	0.36	0.051
MO_Dearborn DS	2019	October	217	1.2	0	0.05	0.004
MO_Hardy Creek	2019	October	203	0.8	0.02	0.3	0.045
MO_Cascade	2019	October	203	1.2	0.11	0.23	0.038
MO_LPPC US	2020	May	222	2	0	0.15	0.014
MO_LPPC DS	2020	May	145	16	0.03	0.16	0.013
MO_Craig	2020	May	223	2	0	0.16	0.006
MO_Dearborn US	2020	May	211	4	0	0.16	0.015
MO_Dearborn DS	2020	May	215	4	0	0.15	0.015
MO_Hardy Creek	2020	May	217	0	0	0.17	0.009
MO_Cascade	2020	May	218	15	0	0.16	0.012
MO_LPPC US	2020	July	211	0	0.04	0.26	0.009
MO_LPPC DS	2020	July	195	1	0.10	0.21	0.018
MO_Craig	2020	July	210	0	0.03	0.25	0.013
MO_Dearborn US	2020	July	202	4	0.03	0.27	0.005
MO_Dearborn DS	2020	July	204	2	0.03	0.25	0.009
MO_Hardy Creek	2020	July	208	0	0.03	0.25	0.015
MO_Cascade	2020	July	207	4	0	0.28	0.017
MO_LPPC US	2020	October	189	2	0.07	0.32	0.039
MO_LPPC DS	2020	October	214	3	0.05	0.14	0.008
MO_Craig	2020	October	193	3	0.04	0.3	0.037
MO_Dearborn US	2020	October	185	2	0.05	0.28	0.035
MO_Dearborn DS	2020	October	187	1	0.04	0.27	0.03
MO_Hardy Creek	2020	October	186	0	0.02	0.25	0.042
MO_Cascade	2020	October	185	1	0	0.23	0.026

Sample Site	Year	Sample Period	TDS (mg/L)	TSS (mg/L)	Inorganic Nitrogen (mg/L)	Total Nitrogen (mg/L)	Total Phosphorus (mg/L)
MO_LPPC US	2021	May	235	2.9	0.01	0.24	0.019
MO_LPPC DS	2021	May	143	26.0	0.07	0.2	0.041
MO_upstream Craig	2021	May	237	3.8	0.01	0.26	0.023
MO_Dearborn US	2021	May	223	4.0	0.02	0.26	0.023
MO_Dearborn DS	2021	May	215	6.0	0.02	0.25	0.023
MO_Hardy Creek	2021	May	219	5.0	0.01	0.48	0.023
MO_Cascade	2021	May	222	7.0	ND	0.24	0.029
MO_LPPC US	2021	July	224	2.9	ND	0.23	0.023
MO_LPPC DS	2021	July	219	1.6	ND	0.18	0.02
MO_upstream Craig	2021	July	230	5	ND	0.23	0.04
MO_Dearborn US	2021	July	225	1.5	ND	0.21	0.019
MO_Dearborn DS	2021	July	232	1.6	ND	0.21	0.017
MO_Hardy Creek	2021	July	229	1.8	ND	0.21	0.015
MO_Cascade	2021	July	222	2.1	ND	0.22	0.019
MO_LPPC US	2021	Sept	228	5	0.02	0.33	0.042
MO_LPPC DS	2021	Sept	217	4	0.02	0.22	0.025
MO_upstream Craig	2021	Sept	227	3.5	ND	0.33	0.039
MO_Dearborn US	2021	Sept	229	2.9	ND	0.28	0.035
MO_Dearborn DS	2021	Sept	228	3.0	ND	0.28	0.034
MO_Hardy Creek	2021	Sept	226	6	ND	0.29	0.040
MO_Cascade	2021	Sept	230	1.9	ND	0.24	0.022
			Hauser Reservoir	Holter Reservoir			
PPC trip_Spring Creek	2021	July	508	8	0.18	0.31	0.045
Canyon Ferry Reservoir							
MO trib_Sixteenmile Cr.	2021	July	292	8	ND	0.21	0.012
MO trib_Big Springs Ditch	2021	July	248	5	0.25	0.32	0.021
MO trib_Dry Creek	2021	July	215	15	0.15	0.57	0.056
MO trib_Greyson Cr.	2021	July	196	104	0.01	0.41	0.157
MO trib_Deep Cr.	2021	July	219	33	ND	0.20	0.041
MO_Townsend	2021	July	210	9	0.07	0.35	0.04
MO trib_Magpie Cr.	2021	July	426	16	ND	0.07	0.013